

Successful Blossom Thinning and Crop Load Regulation for Organic Apple Growing with Potassium-bi-carbonate (Armicarb®): Results of Field Experiments over 3 Years with 11 Cultivars

F. P. Weibel¹⁾, B. Lemcke¹⁾, U. Monzelio¹⁾, I. Giordano¹⁾ and B. Kloss²⁾

¹⁾Research Institute of Organic Agriculture (FiBL), Frick, Switzerland and ²⁾Master Student 2009 (Dept. Horticulture; Prof. J. Wünsche), University Stuttgart-Hohenheim, Germany)

Summary

With field trials over 3 years in a commercial organic orchard in Switzerland we have tested the efficacy of Armicarb® (potassium-bi-carbonate) for flower thinning in organic apple production. Over time, Armicarb was tested on 11 cultivars, at different application periods, in different concentrations, and always in comparison to other agents that are already allowed for thinning in organic fruit production in the European Union as e.g. lime sulphur, molasses, mechanical rope-thinner or combinations of methods. Armicarb proved to be an efficient and reliable thinning agent with an

efficacy similar to the now recommended methods with rope device, molasses or lime sulphur but has the advantage to be an environmentally very friendly product. On the other hand, the risk for fruit russetting is comparably elevated especially with cultivars 'Elstar', 'Golden Delicious' and 'Gala'. Finally, we have elaborated cultivar-specific recommendations for the use of Armicarb for thinning purposes, which were the basis for the Swiss Federal approval to use Armicarb for thinning in conventional and organic apple production in 2011/2012.

Key words. apple – organic – thinning – crop regulation – potassium-bi-carbonate

Introduction

One of the main challenges in organic apple growing is the regulation of the crop load to, i) prevent bi-annual bearing, ii) improve fruit quality, and iii) save labour costs for manual thinning. Up to now, there are only few methods and agents allowed for certified organic agriculture: e.g. the mechanical rope thinner device (BERTSCHINGER et al. 1998, WEIBEL and WALTHER 2003, LAFER 2009, KONG et al. 2009). After WEIBEL et al. 2008, however, 1–2 treatments with the rope thinner alone seldom provide a satisfying result and should be combined with a desiccant agent such as e.g. molasses. Also with 2–3 molasses treatments during flowering period, for rewarding results a combination with the rope device is recommended by the latter authors. In most EU countries and in the US lime sulphur is the standard thinning agent. At dosages of 2–2.5 vol.% and 2–3 treatments over flowering period it provides a fairly good efficacy, and induces little risk for fruit russetting. In Switzerland, however, lime sulphur is not registered by the Federal authorities because of its potential user toxicity.

For the use as a contact fungicide Armicarb (potassium bi-carbonate; KHCO_3) is already licensed for organic apple

production. The active component is 85 % potassium-bi-carbonate, which acts on fungi by changing the pH and the osmotic pressure plus the direct ionic effect of potassium-bi-carbonate on the cell walls (Stähler Suisse SA, Zofingen, CH). After promising pre-trials in 2006 and 2007 to apply Armicarb also for crop regulation, we conducted from 2009–2010 replicated thinning trials with several cultivars under commercial orchard conditions with applications on larger plots and using mostly a commercial orchard sprayer. The main questions to answer were: i) thinning effect of Armicarb in comparison to other organic methods; ii) thinning effect and negative side effects (e.g. phytotoxicity) of Armicarb with different cultivars; iii) optimal concentration for different cultivars; iv) optimal application period and frequency. For this publication, we describe and discuss mainly the results related to Armicarb.

Material and Methods

The field trials were conducted on the commercial organic fruit farm of family Ch. Vogt at Remigen. Situated at the edge of the eastern Swiss Jura Mountains, 450 m above sea level; av. temp. $9.1\text{ }^\circ\text{C y}^{-1}$; av. rainfall 900 mm y^{-1} ;

soil is a pseudo-gleyic, medium-deep brown soil of 18.9 % clay, 45.8 % silt and 35.1 % sand, pH(H₂O) is 5.5–5.9.

The experiments were performed in 2008, 2009 and 2010, and the last return-to-bloom assessment was in April 2011. Treatments tested are described in Table 1. Per cultivar, there were usually 3, minimum 2 replicated plots randomly distributed. A plot consisted of usually 18–27 trees (minimum 9). Most trees were on rootstock M9 (except 'Maigold' on M27, and 'Topaz' on M27 in 2009); in full production age between 7 and 18 years old; spaced 1 × 3 m and under a hail protection net. Usually 10, minimum 5 representative trees per plot were selected as measuring and counting trees. Flowering intensity of the trees used for the experiment was at least 75 % but mainly 90–100 %. Usually the products were applied with a commercial orchard sprayer (Lochmann RPS) using 1000 L water per ha; case-wise a motor backpack sprayer (Birchmeier M155) was used. Usually the test agents were applied twice during flowering stage F and F₂ (BBCH 61–65); 3 applications were occasionally necessary when flowering period was long. The rope device was applied at pre-bloom at stage red tip (E, BBCH 57) at high driving speed (9–11 km h⁻¹) in order to keep physical damages to leaves and branches as low as possible. Because the agents tested (Table 1 Arnicarb treatments, Table 2 other agents and methods) are all desiccants, we applied them at warm days (around 20 °C at midday) with no rain announced for the following 24 h, and when a maximum of un-pollinated flowers were open, thus, spraying time began from 9–10 a.m. on. For the rope device, however, we aimed for colder, cloudy weather to enhance the physiological shock of the treatment (according to WEIBEL et al. 2008). The shadow net was mounted, according to the recommendations of KOCKEROLS et al. (2008) and WIDMER et al. (2009) in average 25 days after full bloom; then fruit diameters reached

8–10 mm, thus timing for shading was also comparable to the studies of GREENE and GROOME (2010) and LAFER (2008).

Fruit set was counted before and after June drop: from each measuring tree 4–6 representative branches in the centre zone of the canopy were chosen. Over the entire length of each branch the amount of fruit clusters and the number of fruits per cluster (0, 1, 2, 3+) was counted using a multiple hand counter tally. Russetting was assessed as % incidence of affected fruits; when treatment-induced russetting seemed to be more intensive, we also assessed severity as percentage of affected fruit skin. Few days before harvest final fruit set was estimated as percentage from an optimal crop load (set as 100 %) and 25 representative fruits per replicated plot were collected to assess fruit diameter and weight of the fruits (in these On-Farm trials yield per tree assessments were not possible to carry out, and the focus of the study was fully on the relative thinning effect of the different treatments). To assess the treatments' influence on bi-annual bearing, return to bloom was counted in mid-April of the following year as percentage of flower buds of total buds.

For statistical analysis we used ANOVA models (treatment, cultivar, replicated block (nested with cultivar) and interaction cultivar*treatment). For multiple treatment comparison a post ANOVA Tukey test was performed ($p < 0.05$; JMP V. 8.0.1, SAS Inc.).

Results

Experiments in 2008

Fig. 1 shows example-wise the thinning effects counted before June drop in 2008 on cv. 'Elstar'. The treatments

Table 1. Variants tested with Arnicarb from 2008–2010 (beside untreated control and hand thinning of 2/3 of the flower clusters).

| Year | Application frequency and timing | Concentrations (kg ha ⁻¹) | Cultivars |
|------|---|---------------------------------------|---|
| 2008 | 3 × during flowering period (F–F ₂ , BBCH 61–65) | 5, 10, 15, 20 | 'Golden Del.', 'Idared', 'Elstar', 'Maigold' |
| 2009 | 1 × at F(61) or F ₂ (63) or F ₂ (64) or F ₂ (65) or at T-stage | 20, 15 at T-stage | 'Topaz*', 'Otava*' |
| | 2 × at F(61) and F ₂ (65) | 10, 15, 20 | 'Golden Del.', 'Gala', 'Elstar', 'Maigold', 'Topaz*' (M9), 'Topaz*' (M27), 'Otava*' |
| 2010 | 2 × at F(61) and F ₂ (65) | 15 | 'Golden Del.', 'Braeburn', 'Pinova', 'Gravensteiner', 'Topaz*', 'Otava*', 'Ariane*' |
| | 2 × at F(61) and F ₂ (65) | 20 | 'Maigold' |
| | 2 × at F(61) and F ₂ (65) | 15 | 'Elstar', 'Topaz*' |
| | combined with rope thinner | | |

* scab resistant cultivar

Table 2. Treatments tested in comparison to Armicarb from 2008–2010 (beside untreated control and hand thinning of 2/3 of the flower clusters)

| Product | Description | Application |
|---|---|---|
| Rope device | “Gessler” (Friedrichshafen, DE), 286 Nylon ropes of 50 cm length on a 2 m vertical axis with 300 rev. min ⁻¹ . | 1–2 at stage (E, BBCH 57) at 9–11 km h ⁻¹ driving speed |
| Covering with Shadow net | “AGROFLOR” (Nendeln, FL) with 74 % light reduction (together with hail nets reduction up to 90 % possible) | Covering for 3–5 days, 20 days after full bloom (J, BBCH 74) |
| Lime sulphur | Ca-Polysulphid 381 g L ⁻¹ “Polisenio”, IT | 2.5 vol%; 2–3 × during flowering period (F–F ₂ , BBCH 61–65) |
| Vinasse Also in combination with rope device application (see above) | Molasses from sugar beet “Bioorga-NK-flüssig” (60 g N L ⁻¹ , 70 g K L ⁻¹); Hauert HGB Dünger AG, Switzerland | 5–7 vol.%; 2–3 × during flowering period (F–F ₂ , BBCH 61–65) |
| Acetic Acid | “Apfelessig” for cooking purposes with 5 g AA L ⁻¹ (Bio Farm, Switzerland) | 3 vol%; 2–3 × during flowering period (F–F ₂ , BBCH 61–65) |
| “Black oil” | Self made mixture of pine oil (NuFilm 1 ml L ⁻¹) and dust of active carbon (25 g L ⁻¹) to induce a micro-shading of the flower clusters by the black colour | 2–3 × during flowering period (F–F ₂ , BBCH 61–65) |
| Goemar® | An algae substrate containing natural GA 14 and micro-nutrients, Stähler Suisse SA, Zofingen, CH | 0.3 vol.% 2–3 times during flowering period (F–F ₂ , BBCH 61–65) |

effects were similar but less expressed in the parallel trials on cv. ‘Idared’. Armicarb (in that year at a dosage of 20 kg ha⁻¹) had a strong, in this case almost too radical thinning effect by decreasing the fruit set from 159 fruits per 100 flower clusters (FICI) to 49. This corresponds to a thinning effect of 69.4 %. Like this, the Armicarb treated trees had only half of the fruit set compared to hand thinning (removal of 2/3 of the flower clusters) and the organic standard treatment rope-device plus vinasse, both showing thinning effects of 33 %. Net-shadowing caused a far too intensive fruit drop down to only 16 fruits 100 FICI⁻¹ remaining.

The effect of the treatments on crop load before harvest in 2008 can be seen in Fig. 2: the untreated control was clearly over loaded with 173.3 % of an optimal crop load; Armicarb treated trees were slightly over-thinned showing 91.7 % of an optimal crop load, shadow nets clearly over-thinned to only 51.7 %, meanwhile the positive control treatments like hand thinning and rope device plus vinasse were between 112 and 123 %; Goemar and Acetic Acid had no effect.

As a consequence, fruit weight of Armicarb treated trees increased by 25.2 % from 130 g fruit⁻¹ in the untreated control to 174.3 g fruit⁻¹. Return to bloom in the Armicarb treated ‘Elstar’ plots in the following year with 89.6 % flower buds was clearly higher, almost too high, com-

pared to hand thinning and rope device plus vinasse with only 7–23 % flowering buds whereas untreated control developed only 0.9 % of flower buds (data per treatment not shown in detail).

The trials with different Armicarb concentrations in 2008 with cv. ‘Golden Delicious’, ‘Idared’, ‘Elstar’ and ‘Maigold’ revealed that concentrations must be between 10–20 kg ha⁻¹. The results showed a tendency that with most cultivars 5 kg have almost no thinning effect, 15 kg ha⁻¹ are significantly more effective than 10 kg ha⁻¹, but 20 kg ha⁻¹ do not further improve the thinning effect compared to the 15 kg ha⁻¹ concentration. Furthermore, the incidence of fruit russeting – mainly with ‘Elstar’ – increased at 15 and 20 kg ha⁻¹ (data not shown in detail, see results 2009).

Experiments in 2009

In 2009 the experiments on different Armicarb concentrations were repeated with 10, 15 and 20 kg ha⁻¹ on cv. ‘Gala’, ‘Golden Del.’, ‘Elstar’, ‘Maigold’, ‘Topaz’ (on M9, scab resistant), ‘Topaz’ (M27) and ‘Otava’ (scab resistant). The results confirmed that a relevant thinning effect can be achieved only from 15 kg ha⁻¹ on. This trend was obvious for all cultivars tested. In that year, the thinning effect by 15 kg ha⁻¹ Armicarb as it was assessed before June




Fig. 1. Number of fruits/100 clusters with cv. 'Elstar' under 8 different thinning treatments counted before June drop in 2008. Represented are Box Plots with the great mean line (horizontal line). AZE = acetic acid. Vin = vinasse, LSD = least significant difference.

drop was around 40 % with 'Topaz' on M9 and M27 and 'Elstar', around 30 % with 'Maigold' and 'Otava' but only 2 % with 'Golden Del.' (data not shown). After June drop (Fig. 3), 'Golden Del.' 'caught up' to a thinning effect of 30 %, similar to 'Elstar', 'Gala' and 'Topaz' (on M9 and M27); with 'Maigold', due to a high natural June drop, only a 9.4 % thinning effect resulted at this date. With most cultivars the 15 kg ha⁻¹ Arnicarb concentration led to a close to optimal crop load before harvest. Exceptions were 'Maigold', where a concentration of 20 kg ha⁻¹ gave a better final result without a concerning increase of russetting. In this year, especially with 'Elstar', 'Gala' and 'Golden Del.' the 15 kg ha⁻¹ dosage of Arnicarb increased the incidence of fruits affected with russetting in a magnitude of 10–17 %.

The data on return to bloom as percentage of flower buds in the following year (2010) did not reveal significant treatment effects except for 'Otava' with an increase of 61 % flower set. Nevertheless, in the plots treated with 15 kg ha⁻¹ Arnicarb, crop load before harvest in 2010 was improved towards optimal fruit set in the magnitude of 9 % ('Elstar' and 'Otava') to 34 % ('Topaz') (data not shown in detail).

In the separate trial to test different timing of Arnicarb with cv. 'Topaz' and 'Otava' with only a single application, we could see that the thinning effect of later Arnicarb applications at stage F₂ 65 is superior (22 %) than with earlier applications at stage F₂ 61 or 62 or 63 (8.2 %). The reason for this pattern is that the later the more flowers are open and affected by the agent. Also the incidence of russetting increased with later applications

from 4 % at F₂ 62 up to 10 % at F₂ 65. Russetting damages were particularly severe – reaching 22 % incidence – in the case where 20 kg ha⁻¹ Arnicarb were applied on 'Topaz' at late flowering stage F₂ 65 shortly before it began to drizzle with rain. We assume that under these circumstances Arnicarb got entirely in solution and too intensively into contact with the fruit epidermis. The data of the timing trial are not shown in detail.

In 2009 also different alternative methods were tested on cv. 'Topaz' and 'Otava'. Natural fruit fall, however, was high this year due to a relatively cold climate causing sub-optimal conditions for assimilation for the trees during May. For this reason, even in the untreated control variant and with both cultivars, crop load at harvest was only 10 % too high in the untreated control plots. At the fruit counting date before June drop, the treatments 2 × 20 kg ha⁻¹ Arnicarb and rope device plus 3 × 7 % vinasse showed a significant but low thinning effect of 12.8 and 18.9 %. 3 × 2.5 % lime sulphur with 30.4 % thinning effect was significantly more effective than the latter treatments. As in 2008, net shadowing reduced fruit set too radically by 67.1 %. The interaction treatment * cultivar was not significant. The results of this trial are not shown in detail.

Experiments 2010

In 2010 Arnicarb concentration trials were performed with 15 kg ha⁻¹ on cv. 'Golden Del.', 'Braeburn', 'Pinova', 'Gravensteiner', 'Topaz', 'Otava', 'Ariane' (scab resistant), and with 20 kg ha⁻¹ on 'Maigold'. The good thinning




Fig. 2. Percentage of optimal fruit load (visually estimated) before harvest with cv. 'Elstar'. Dotted line = optimal fruit load (100 %). Vin = vinasse, AZE = acetic acid. LSD = least significant difference.

effect of Armicarb at these concentrations could be confirmed, though, as a consequence of the varying intensity of the natural June drop in the control plots, the thinning effect (expressed as % difference to the untreated control) could vary considerably between the fruit counting before and after June drop (Table 3). Nevertheless, with all cultivars, either before or after June drop significant and for practical fruit growing relevant thinning effects in the magnitude of 13 % ('Otava') to 52 % ('Gravensteiner') could be achieved.

Fig. 4 shows that beside the total thinning effect, with all cultivars, the 15 kg ha⁻¹ Armicarb treatment also reduced the proportion of flower clusters with 3 or more and 2 fruitlets in favour of clusters with 1 or 2 fruitlets, respectively; from the perspective of the fruit grower a most desirable pattern.

The 2010 trials to compare different alternative methods carried out on cv. 'Elstar' and 'Topaz' included applications of lime sulphur, Armicarb standard (15 kg ha⁻¹), Armicarb 15 kg ha⁻¹ in combination with a rope device application at stage E (59) and "Black Oil". In 2010 the weather conditions right after blooming period were unusually cold and rainy during 24 days. The conditions for assimilation and fruit set were therefore sub-optimal. For these reasons, the results of these method comparison trials are to some extent difficult to interpret and not shown in detail: The counting before June drop revealed a generally very high fruit set with 230 to 245 fruits 100 FICl⁻¹ over all treatments including the by 2/3 hand thinned trees (as a compensation reaction, these latter trees kept most of the remaining fruits and thus had a

high number of > 2 and > 3 fruits per fruit cluster). At this date, only the lime sulphur treatment (2 × 2.5 vol.%) revealed a moderate thinning effect of 22.3 %. After June drop, however, fruit set dropped drastically to 85–111 fruits 100 FICl⁻¹ in all treatments (including untreated control) with the exception of the lime sulphur treatment, which decreased to 58 fruits 100 FICl⁻¹, and therefore was even over-thinned. Consequently at harvest 2010, the remaining treatments appeared with a near optimal crop load, again without significant treatment differences.

In the trials of 2010 no variant showed signs of treatment-induced russeting. For this reason, the assessments of russeting incidence and severity were not carried out that year.

Discussion

Armicarb is a well-known product against scab and sooty blotch on apple (TAMM et al. 2006). Since presently, no active compound is allowed for fruit thinning in organic apple production in Switzerland, Armicarb is an interesting candidate for organic apple thinning. PFEIFFER (2008) studied the thinning effect of Armicarb where it was applied as a fungicide at a rate of 5 kg ha⁻¹: Very much in line with our study, there was too little thinning effect of Armicarb at that dosage. To our knowledge, so far there are no other longer-term and multi-cultivar studies published (including return to bloom data) where Armicarb was tested at higher dosages as a thinning agent. In our experiments, we could show that Armicarb has a high thinning




Fig. 3. Armicarb concentration trials in 2009 with cv. 'Elstar', 'Gala', 'Golden Del.', 'Maigold', 'Otava', 'Topaz' (on rootstocks M9 and M27) with 0, 10, 15 and 20 kg ha⁻¹ Armicarb in 1000L water 2 × during bloom: Number of fruits per 100 flower clusters (FrCl) after June drop; thinning effect compared to control (%); crop load before harvest (% of optimum); incidence of fruit russetting (% incidence); fruit weight (g). Interaction treatment * cultivar is significant; in all cases differences between control and 15 kg ha⁻¹ were significant, except for 'Maigold' where this is the case with 20 kg ha⁻¹.

potential and that its effect is – to a manageable extent – cultivar and concentration dependent. Our concentration experiments over 3 years showed that Armicarb has a significant thinning effect with a concentration of 15 kg ha⁻¹ for 10 of the cultivars tested and with 20 kg ha⁻¹ for cv. 'Maigold'. Interestingly, a higher concentration did not induce a stronger thinning effect but increased the incidence of russetting. Using Armicarb as thinning method reduces labour costs compared to hand thinning and showed better results than the previously best method (under Swiss conditions) which was the combination of rope thinner followed by 2–3 applications with vinasse. Moreover Armicarb, related to baking powder, is considered as very eco-friendly substance.

Our results are in line with previous findings (WEIBEL et al. 2008; LAFER 2009) confirming that the rope device combined with a desiccant as vinasse or lime sulphur is a fairly efficient thinning method. Furthermore, with a vinasse concentration of 5–7.5 % and 2–3 applications during blooming phase, we did not observe phytotoxic effects. To avoid damages on wood, spurs and leaves (BAAB and LAFER 2005) and a too intense physiological shock of the tree (mechanism described by UNTIEDT and BLANKE 2001) it is important to use the rope device as softly as possible by driving at high tractor speed (9–11 km/h) and at moderate rev./min. of the rope spindle (WEIBEL and WALTHER 2003). For orchards with lager trees than slim spindles, the rope machine type "Bonn" with 3 horizontal rope axes

Table 3. Thinning effect of Armicarb ($2 \times 15 \text{ kg ha}^{-1}$) in the 2010 trials with different cultivars as assessed before and after June drop. Asterisks indicate statistically significant effects (post ANOVA Tukey HSD tests at $p < 0.05$).

| Cultivar | Thinning effect before June drop (%) | Thinning effect after June drop (%) |
|---------------------------------------|--------------------------------------|-------------------------------------|
| 'Braeburn' | 22.4* | 21.8* |
| 'Golden Del.' | 30.0* | 44.5* |
| 'Gravensteiner' | 29.0* | 52.3* |
| 'Otava' | 11.9* | 13.2* |
| 'Pinova' | 24.6* | 15.6 |
| 'Topaz' | 9.8 | 37.1* |
| 'Maigold' (20 kg ha^{-1}) | 45.8* | 16.7 |

can reach better also the inner tree parts (DAMEROW et al. 2007, KONG et al. 2009).

In our experiment, hand thinning by 2/3 was consequently included as 'positive' control treatment (DENNIS 2000 and 2002). However, beside unaffordable costs for labour, hand thinned trees tend to compensate the removed flowers by keeping a very high proportion of fruits on the remaining flower clusters, which finally results in a high proportion of flower clusters with 2 and 3 and more fruits (KLOSS and WEIBEL 2009, Bachelor Thesis University Hohenheim).

Goemar® GA14, vinegar (5 g L^{-1} acetic acid) and "Black Oil" did not show significant thinning effects in these trials and are therefore not profoundly discussed.

Our results on shadowing are in line with previous studies, which showed that tree shading is an efficient, thinning method (MCARTNEY et al. 2004, STADLER et al. 2005, KOCKEROLS et al. 2008). According to STADLER et al. (2005) we mounted the nets 22–23 days after full bloom for 5–6 days for 'Elstar' and 4 days for 'Idared'. However this resulted in clear over-thinning with both cultivars in both years. We suppose that in our experiment, this shading duration was too long under the circumstances given with low natural radiation (MCARTNEY et al. 2004) and an additional hail net.

Conclusions

From our experiments, and for the cultivars and conditions tested, we draw the following conclusions for the practical application for organic thinning measures during bloom:

1. With the majority of cultivars, 2 applications during bloom of 15 kg ha^{-1} potassium-bi-carbonate (Armicarb®) gave a satisfying result. Only with cultivar 'Maigold' $2 \times 20 \text{ kg}$ gave a better result.
2. When applying Armicarb on not yet tested cultivars, 15 kg ha^{-1} is a recommendable starting concentration for tests. The optimum concentration, however, can range between $12\text{--}20 \text{ kg ha}^{-1}$ depending on cultivar, flower set and climatic conditions.




Fig. 4. Mean numbers of fruits per flower cluster per cultivar and treatment, counted before June drop 2010 in the treatment with $2 \times 15 \text{ kg ha}^{-1}$ Armicarb; 'Maigold' with 20 kg ha^{-1} Armicarb.

3. The application of Armicarb should take place at warm, sunny days without rain in the following 12 h and at a time with a maximum of still un-pollinated flowers wide open. Thus, depending on the duration and intensity of the blooming phase, 2–3 applications are necessary.
4. With some cultivars like ‘Elstar’, ‘Gala’, ‘Golden Delicious’ and climatic conditions that favour russetting, the use of Armicarb for thinning can increase incidence and severity of fruit russetting. In particular, it has to be avoided that it starts drizzling shortly after the application of Armicarb.
5. We achieved good thinning results also with 2–3 applications of lime sulphur at 2–2.5 vol.%, and vinasse at 5–7 % concentration at the same conditions as mentioned above for Armicarb. Vinasse is particularly efficient with e.g. cultivar ‘Topaz’.
6. The efficacy of vinasse can or even should be improved with a gentle application of the rope thinning device at stage red tips (E/59), followed by two or three applications of the desiccant agent during flowering period.
7. Rather than by the physical removal of the flower organs, the thinning effect of the rope device should be the consequence of a physiological shock followed by a lack of assimilates for the development of the fruitlets and formation of ethylene (UNTIEDT and BLANKE 2001, GREENE 2002, WÜNSCHE and FERGUSON 2005). Thus, this method should be applied at colder, cloudy days to increase the effect of the photosynthesis decreasing effect. The Rope device should not be used during full bloom because then also many primary leaves will be destroyed, and this harsh physiological shock will cause long-lasting negative effects (e.g. compensatory shoot growth in summer etc.).
8. Shadowing has a proven strong thinning effect. However, in our experiments we provoked over-thinning in spite of having applied the method as recommended. The Swiss organic fruit growers having tried the method on a larger scale gave feedback that for them shadowing is too labour intensive, and also had too varying results with over- and underthinning.

Acknowledgements

Many thanks go to the family Vogt, especially to Christian, who allowed us to install the trials in their orchards with great trust, and who very carefully and reliably applied the Armicarb, vinasse and rope device variants with his professional equipment. A special thank goes to Albert Widmer from Agroscope Changins-Wädenswil (ACW) for borrowing us the shadow nets and for his advice on shadowing. The Stähler Swiss Company co-financed the study and provided us with the products Armicarb and Goemar for our above large trials. A special thank goes to Stähler’s staff Matthias Refardt, Thomas Steiner and David Vullie-

min for their most appreciated support with counting the fruit clusters.

References

- BAAB, G. and G. LAFER 2005: Kernobst. Leopoldsdorf, Austria: Österreichischer Agrarverlag Druck- und Verlags-GmbH.
- BERTSCHINGER, L., W. STADLER and F. P. WEIBEL 1998: New methods for an environmentally safe regulation of flower and fruits set and alternate bearing of apple crop. *Acta Hort.* **466**, 64–70.
- DAMEROW, L., M. BLANKE and P. SCHULZE-LAMMERS 2007: [Crop load in pome fruit cultivation] Regulierung der Fruchtbehangsdichte im Kernobstanbau. Forschungsbericht Nr. 143. Rheinische Friedrich-Wilhelms-Universität Bonn, Landwirtschaftliche Fakultät (In German).
- DENNIS, F.G. Jr. 2000: The history of fruit thinning. *Plant growth regulation*, **31**, 1–16.
- DENNIS, F.G. Jr. 2002: Mechanism of action of apple thinning chemicals. *Hortscience* **37**, 471.
- GREENE, D.W. 2002: Chemicals, timing and environmental factors involved in thinner efficacy on apple. *Hortscience* **37**, 447–448.
- GREENE, D. W. and P. GROOME 2010: Effect of thinner and time shading on fruit set of McIntosh apples. *Acta Hort.* **884**, ISHS 2010; 505–510.
- KOCKEROLS, K., A. WIDMER, M. GÖLLES, L. BERTSCHINGER and S. SCHWAN 2008: Ausdünnung von Äpfeln durch Beschattung. *AGRARForschung* **15**, 258–263 (In German).
- KONG, T.; L. Damerow and M. BLANKE 2009: [Influence of selective mechanical thinning methods on ethylensynthesis, yield and fruit quality on pome fruit] Einfluß selektiver mechanischer Fruchtbehangsregulierung auf Ethylensynthese, Ertrag und Fruchtqualität bei Kernobst. *Erwerbs-Obstbau (Springer Heidelberg)* **51**(2), 39–53 (In German).
- LA FER, G. 2008: [Thinning by Shading Methods] Ausdünnung durch Beschattung eine alternative Methode zur Ertragsregulierung. *Obstbau*, **4**, 229–230 (In German).
- LA FER, G. 2009: [Experiences with Mechanical Thinning in Styrian, Organic Fruit Cultivation] Erfahrungen mit der mechanischen Ausdünnung im steirischen Bioobstbau. *Gleisberger Bioobstbautage 2009* (In German).
- M CARTNEY, S., M. WHITE, I. LATTER and J. CAMPBELL 2004: Individual and combined effects of shading and thinning chemicals on abscission and dry-matter accumulation of ‘Royal Gala’ apple fruit. *J. Hort. Sci. Biotech.* **79**, 441–448.
- PFEIFFER, B. 2008: Results from scab trials with Armicarb in the years 2006 and 2007. In: “Proceedings 12th International Conference on Cultivation Techniques and Phytopathological Problems in organic Fruit Growing”; pp. 41–46, ed. Fördergemeinschaft Ökologischer Obstbau e. V.: Weinsberg, Germany.

- STADLER, W., A. WIDMER, E. DOLEGA, M. SCHAFFNER and L. BERTSCHINGER 2005: Fruchtausdünnung durch Beschattung der Apfelbäume – eine Methode mit Zukunft? Schweizerische Zeitschrift für Obst- und Weinbau **10**, 10–13 (In German).
- TAMM, L., T. AMSLER, H. SCHÄRER and M. REFARD 2006: In: Proceedings 12th International Conference on Cultivation Techniques and Phytopathological Problems in organic Fruit Growing (ed. Fördergemeinschaft Ökologischer Obstbau e. V.: Weinsberg, Germany), pp. 87–92.
- WEIBEL, F., V.S. CHEVILLAT, E. RIOS, J.L. TSCHABOLD and W. STADLER 2008: Fruits thinning in organic apple growing with optimized strategies including natural spray products and rope-device. *Europ. J. Hort. Sci.* **73**, 145–154.
- WEIBEL, F. and A. WALTHER 2003: Ausdünnung beim Apfel: beim Fadengerät lässt sich noch einiges herausholen! Proceeding of the Swiss Organic Fruit Symposium, January 3, 2003 at Frick, pp. 36–40 (In German).
- WIDMER, A., K. KOCKEROLS and M. GÖLLES 2009: [Thinning of apples by shading methods] Ausdünnung von Äpfeln durch Beschattung; in Vol. 1, 10. Wissenschaftstagung ökologischer Landbau ETH Zürich 11.–13. Feb. 2009; 141–144 (In German).
- WÜNSCHE, J. N. and B. FERGUSON 2005: Crop Load Interactions in Apple. In JANICK, J. (ed.): *Horticultural Reviews*. Wiley J. und Söhne, Inc., 231–290.

Received 04/10/2012 / Accepted 06/21/2012

Addresses of authors: F.P. Weibel (corresponding author), B. Lemcke, U. Monzelio, I. Giordano, Research Institute of Organic Agriculture (FiBL), Frick, Switzerland and B. Kloss, Master Student 2009 (Dept. Horticulture; Prof. J. Wünsche), University Stuttgart-Hohenheim, Germany, e-mail (corresponding author): franco.weibel@fibl.org.